

Overall **N**utritional **Q**uality **I**ndex
Version 1
(ONQI.v1)

- REFERENCE MANUAL -

Prevention Research Center
Yale University School of Medicine

Griffin Hospital
Derby, CT

Copyright © 2007 Griffin Hospital

Authors:

David L. Katz, MD, MPH, FACPM, FACP

Associate Professor, adjunct, Public Health

Director, Prevention Research Center

Yale University School of Medicine

www.davidkatzmd.com

Project Director

Valentine Yanchou Njike, MD, MPH

Data Analyst/Programmer

Debbie Kennedy, PhD

Zubaida Faridi, MD, MPH

Project Coordinators

Judy Treu, MS, RD

Lauren Q. Rhee, MS, RD

Research Dietitians

This project was fully supported by Griffin Hospital, Derby, CT.

Primary Contact:

David L. Katz, MD, MPH, FACPM, FACP

Yale University Prevention Research Center

Griffin Hospital, 2nd Floor

130 Division St.

Derby, CT 06418

www.davidkatzmd.com

Phone: (203) 732-7194

Fax: (203) 732-1264

Email: Katzdl@pol.net

david.katz@yale.edu

Administrative Assistant, Helen Day

helen.day@yalegriffinprc.org

Précis

The United States is the epicenter of a global obesity pandemic. Insulin resistance, metabolic syndrome, and type 2 diabetes are epidemic as well. What used to be “adult onset” diabetes is diagnosed with increasing regularity in children under the age of ten. The combination of dietary patterns at odds with recommendations and relative lack of physical activity is, or soon will be, the leading cause of premature death in the land.

There is widespread consensus among experts that improvements in activity levels and dietary patterns have enormous potential to promote health and prevent disease. Healthful eating and activity patterns (along with other sensible behaviors, such as tobacco avoidance) could reduce the rate of heart disease by as much as 80%, diabetes by up to 90%, and cancer by nearly 60%. Were any such advances the result of a medical breakthrough, it would surely warrant- and earn- a Nobel Prize. As it is, they are the potential products of applying what we already know.

In the case of diet, there are two ways to close the gap between how we eat at present and how we should eat for optimal health. One involves a fundamental shift in the pattern of the diet, as reflected in such advice as “eat more fruits and vegetables.” Such advice is valid, and important, but subject to considerable resistance. At the current rate of progress toward Healthy People 2010 goals for fruit and vegetable intake, for instance, it is only modest hyperbole to project hitting the target by approximately the year 3010!

There is another way to improve dietary patterns and that is one food choice at a time. The range in nutritional quality for every food category represented on supermarket shelves, from greens to granola bars, sandwich meat to salad dressing, cookies to cooking oils, and even the proverbial soup to nuts- is vast. Choosing the most nutritious offerings in each category offers a powerful means to reduce intake of calories, sodium, added sugar, and harmful fats, while increasing intake of fiber, beneficial nutrients, vitamins, and minerals.

But between the average consumer and those choices is a rolling sea of competing claims, half truths, marketing hype, and hopeless confusion.

Upon those troubled waters, the ONQI offers a lighthouse beam. An unfailing, ever reliable guide to better nutrition both within and across food categories. A beacon to better health that comes of better eating. An illumination... of how actually to get there from here.

Select Sources:

Cordain L, Eaton SB, Sebastian A, Mann N, Lindeberg S, Watkins BA, O'Keefe JH, Brand-Miller J. Origins and evolution of the Western diet: health implications for the 21st century. *Am J Clin Nutr.* 2005;81(2):341-354

Dwyer J. Starting down the right path: nutrition connections with chronic diseases of later life. *Am J Clin Nutr.* 2006;83(2):415S-420S

Diet, nutrition and the prevention of chronic diseases. *World Health Organ Tech Rep Ser.* 2003;916:i-viii, 1-149.

Eyre H, Kahn R, Robertson RM; American Cancer Society, the American Diabetes Association, and the American Heart Association. Collaborative Writing Committee. Preventing cancer, cardiovascular disease, and diabetes: a common agenda for the American Cancer Society, the American Diabetes Association, and the American Heart Association. *Diabetes Care.* 2004;27(7):1812-1824.

Hu FB, Willett WC. Optimal diets for prevention of coronary heart disease. *JAMA.* 2002;288(20):2569-2578.

Katz DL. Diet and Health Promotion. In: Katz DL with Friedman RSC. *Nutrition in Clinical Practice.* 2nd ed. Philadelphia, PA: Wolters Kluwer/Lippincott Williams & Wilkins;. In production: October, 2007.

Katz DL. Obesity...be Dammed! What it will take to turn the tide. *Harvard Health Policy Review.* 2006;7:135-151.

Willett WC, Stampfer MJ. Foundations of a Healthy Diet. In: Shils ME, Shike, M, Ross AC, Caballero B, Cousins RJ, eds. *Modern Nutrition in Health & Disease.* 10th ed. Philadelphia, PA: Lippincott Williams & Wilkins;. 2006:1625-1637.

<u>Table of Contents</u>	<u>Page</u>
Title Page	1
Authors	2
Key Contact	3
Précis	4
TOC	6
ONQI Definition / Characterization	8
Mission & Objectives	10
Vision Statement	11
Proviso	12
ONQI Project Overview	13
Origins of the ONQI Project	13
Methods	14
ONQI Scientific Expert Panel	17
ONQI Scientific Advisory Board	19
Informed Contacts (not directly involved)	21
ONQI Presentations	25
ONQI Algorithm	26
Introduction	26
Nutrient Input	27
Determination of Trajectory Scores	32
Threshold Values	33
Trajectory Score Calculation	33
Weighting Coefficients	35
Universal Adjustors	38
Food Categories	42
Categorical Adjustments to Algorithm	44
ONQI Formula	44
Basic Formula	45

<i><u>TOC cont'd</u></i>	<i><u>Page</u></i>
Adjustments to Formula	45
Scores Generated	46
Synopsis	46
Nutrient Database: Populating the ONQI Algorithm	46
Validity Testing	49
Comparison to Other Nutrition Scoring Systems	50
APPENDIX 1: ONQI Program, SAS, with explanatory comments	53
APPENDIX 2: Illustrative Food Rank List	54
APPENDIX 3: Biosketches, Expert Panel & Advisory Board	55
APPENDIX 4: Overview of ONQI On-line Applications	63
APPENDIX 5: Overview of Commercialization Plan	64
APPENDIX 6: Considerations for Future Iterations of the ONQI	65
BIBLIOGRAPHY	67

ONQI Definition / Characterization

The *Overall Nutritional Quality Index* (ONQI) is an algorithm designed to generate a single, summative score for the “overall nutritional quality” of a food based on its micronutrient and macronutrient composition and several other of its nutritional properties (e.g., energy density). The ONQI is further designed to stratify foods into a rank order of relative nutritiousness both universally (i.e., across all food categories) and within specific food categories (e.g., breads, cereals, frozen desserts, etc.), while avoiding the characterization of any food as “good” or “bad” in absolute terms. The ONQI is further designed to convert the pooled knowledge, judgment, and discriminating ability of top experts* in nutrition and its associations with public and personal health into a tool that any consumer can use. Thus, the ONQI enables the “average shopper” to choose foods on the basis of overall nutritional quality with the ease and fidelity of top nutrition experts. By virtue of the ONQI algorithm sophistication and the expansive array of nutrient data it incorporates, the ONQI is applicable to any food item, recipe, meal, meal plan, or overall dietary pattern. It is designed to have applications at point of purchase in retail supermarkets, on food packaging, in restaurants, in print materials (e.g., books, periodicals), and on-line.

***Consulting Participants on the ONQI Scientific Expert / Development Panel (N.B. Not all members of the ONQI development panel remain involved in the project; see advisory board roster, p. 19):**

- Chair: Dr. David L. Katz, Yale University Prevention Research Center
- Dr Keith Ayooob, Albert Einstein College of Medicine
- Dr Leonard Epstein, University at Buffalo, The State University of New York; inventor, Traffic Light Diet
- Dr David Jenkins, University of Toronto; inventor, Glycemic Index
- Dr Francine Kaufman, University of Southern California; Past President, American Diabetes Association
- Dr Robert Kushner, Northwestern University
- Dr Ronald Prior, Arkansas Children's Nutrition Center, USDA HNRC
- Dr Rebecca Reeves, Baylor College of Medicine; Past President, American Dietetic Association
- Dr Barbara Rolls, Pennsylvania State University
- Dr Sachiko St. Jeor, University of Nevada
- Dr John Seffrin, President & CEO, American Cancer Society
- Dr Walter Willett, Harvard University

ONQI Project: Mission & Objectives

- To develop a (the) definitive algorithm to stratify food items within any category, and across food categories, based on overall nutrition quality.
 - *To measure overall nutritional quality using the best science available, encompassing nutritional biochemistry; physiology and metabolism; dietetics; and epidemiology and public health.*

- To develop an algorithm for assessing the overall nutritional quality of foods that entirely avoids the “good food/bad food” dichotomy and controversy.
 - *Specifically, to develop a novel metric for the nutritional quality of individual foods, and the nutrients in those foods, based on the relationship among nutrients in foods, the overall quality of the diet, and associations with health outcomes.*

- To place the ability of top nutrition experts to discriminate among food choices on the basis of nutrition into the hands of every consumer.

- To provide the public a powerful and empowering means to improve dietary intake patterns, and thereby health, one food choice at a time.

ONQI Project Vision Statement:

- A single, trusted, reliable and universal nutrition guidance system available to consumers at every point of purchase, freely accessible on-line, and on every package in the food supply enabling all consumers to choose foods for consumption on the basis of overall nutritional quality with complete confidence, great convenience and ease, and unfailing immunity from any and all potential deception related to marketing.
- An improvement in the approximation of prevailing dietary patterns to recommended dietary patterns, specifically (in the United States) the current version of the *Dietary Guidelines for Americans*.
- An improvement in the public health, as reflected in particular by declining rates of obesity, diabetes, insulin resistance, cardiovascular disease, and cancer, by virtue of improvements in prevailing dietary patterns.

Proviso

The ONQI is a tool, and like any other tool- including the very best- is well suited to a particular function. In the case of the ONQI, that function is to guide food choice/selection on the basis of overall nutritional quality.

Nutritional quality is a measure analogous to density. Just as the density of diamond does not vary with the size of the stone, nor does the nutritiousness of broccoli vary with its serving size. Thus, the ONQI tool in its unadorned application does not offer guidance for portion sizes, or other elements involved in constructing an optimally balanced dietary pattern. To be maximally useful, a tool must be used well, and more or less as intended. The ONQI is no exception.

That said, among the factors considered in scoring the nutritiousness of foods by use of the ONQI are properties that influence satiety. Consequently, foods that score well on the ONQI scale will, in general, facilitate portion control. As for dietary balance, an adaptation of the ONQI applicable to the entire diet- accounting for balance and variety along with the nutritional quality of individual food choices- is under development.

Nonetheless, this delimiting precaution is warranted. A tool used well is empowering; a tool used badly is potentially harmful. A hammer is excellent for driving nails, rather poor for opening windows. The ONQI is a powerful tool, simple to use. But the true measure of its utility, as that of any tool, is contingent upon how it is put to use.

ONQI Project Overview

Origins of the Project

The ONQI project traces its origins to July, 2003. At that time, U.S. Secretary of Health convened a group of 15 academic thought leaders to share ideas for improving dietary intake patterns in the United States, and specifically for curtailing the spread of epidemic obesity, with him, FDA Commissioner Mark McClellan, and the Directors of the NIH & CDC.

The ONQI concept was proposed at that meeting¹. Specifically, the following recommendations were made:

- Develop a universal guidance system based on the relative nutritiousness of foods within and across all categories.
- To do so, convene a multidisciplinary group of leaders in nutrition and its implications of public health under the auspices of the US FDA or the Institute of Medicine (IOM).
- Provide the group the financial support and time required to convert their collective knowledge and judgment into a universally applicable algorithm for nutrition scoring.
- Convert nutrition quality scores into a set of symbols, reflecting relative nutritional quality, interpretable at a glance.
- Apply those symbols to the entire food supply.
- Provide the requisite support for on-going oversight of the scoring system, and its timely revisions based on changes in nutrition science, epidemiology, and/or the food supply.

¹Katz DL. Verbal commentary to Secretary of Health Tommy Thompson, July, 2003, Washington, D.C.

Katz DL. *A food supply for dummies*. Op-Ed: Hartford Courant; NY Newsday; etc. 10/03

While there was strong philosophical support for the recommendation in 2003, and some follow-up discussion with FDA officials responsible for oversight of an advisory group convened to address obesity control, no formal action was taken.

In late 2005, a confluence of circumstances resulted in the willingness of Griffin Hospital, a Yale-affiliated, non-profit community hospital in Derby, CT, and home to Yale University's Prevention Research Center, to provide financial and material support for the very project proposed to the Secretary of Health in 2003.

The ONQI project was thus initiated in late 2005, and the first draft of the ONQI algorithm was created in February, 2006.

Methods

The ONQI project was initiated in an explicit effort to satisfy the criteria proposed to the US Secretary of Health in 2003, and to replicate, to the extent possible, the very process that the FDA or IOM "would have" applied. The salient elements in such a process include:

- Engagement of a multidisciplinary team of nutrition and public health scientists
- Avoidance of conflicts of interest
- Isolation of the science-based development effort from any outside or commercial influence
- Reliance on objective sources of data to the extent possible
- Transparency of methods
- A fully iterative process, with consensus-based decision-making

The ONQI project was implemented in satisfaction of these principles. The sequence of methodologic steps was approximately:

- 1) A review of literature pertaining to nutrition guidance and nutrient profiling from both the US and abroad by staff at the Prevention Research Center.
- 2) Concurrent with #1, a delineation of a master list of potential members of the ONQI Scientific Expert Panel based on a need to include thought leaders in nutrition science and public health nutrition; diverse disciplines; and the representation of leading institutions and organizations.
- 3) A winnowing of the list to achieve a panel of between 10 and 15 participating consultants.
- 4) Invitations extended to a total of 16 nutrition and public health scientists.
- 5) Fourteen of those invited to join the ONQI development panel did so; 3 of these subsequently stepped down. One participant stepped down after recognizing a potential conflict of interest; one did so after recognizing a personal conflict with the goal of the project; and one simply did not participate in the information exchange process and was lost to attrition. The final panel of 11 consultants represented fields from nutritional biochemistry to chronic disease epidemiology, pediatric endocrinology to behavior modification.
- 6) The identities of the consulting scientists were kept in confidence to avoid any and all outside influence.
- 7) The consultants were compensated for their time but did not have any financial interest in the final product; the details of the agreement between the panel members and Griffin Hospital were addressed in a letter of understanding.
- 8) Beginning in early 2006, a draft of the ONQI algorithm and its justifications were circulated to panel members for review and critique. The basic formula is based on the Dietary Reference Intakes (DRIs); Food and Drug Administration (FDA) Nutrition Facts Panel; US Department of Agriculture (USDA) MyPyramid and the Dietary Guidelines for Americans, 2005; and relevant international standards.
- 9) An iterative process of editing took place based on feedback from panel members via e-mail and at regularly scheduled conference calls, with process oversight and logistical support rendered by staff at the Prevention Research Center.

- a. Over a period of 18 months, starting in March 2006, 13 conference calls were held to make revisions to the ONQI based on consensus, and to address validation/testing.
 - b. Work was managed in between calls primarily via e-mail exchanges.
- 10) Once an initial, consensus-based version of the ONQI formula was in circulation, validation testing was initiated. Consensus approval of the form of the algorithm was deemed commensurate with *face validity*.
- 11) Content validity testing was conducted in the following manner:
- a. Members of the ONQI panel were given foods lists of varying length and variety and asked to rank-order the foods on the basis of overall nutritional quality across food categories (lists of approximately 20 foods) and both within and across food categories (lists of approximately 100 foods).
 - b. A mean ranking for the ONQI panel was established.
 - c. The ONQI algorithm was used to rank order the same foods, and correlation analysis was performed.
 - d. This process was terminated with a food list of roughly 100-120 items, and with a correlation coefficient between ONQI and expert panel rankings of between $R=0.88$ and $R = 0.92$, depending on certain fine adjustments to the algorithm.
 - e. At that point, the process was reversed: the ONQI was used to rank order over 1000 foods, and the relevant rank-ordered lists were circulated to the expert panel members for review. Any apparent anomalies identified in the food rankings resulted in scrutiny of the nutrient data entered, and a determination to either acknowledge that the ONQI ranking was correct, or to adjust the algorithm. This process continued until no further adjustments to the algorithm were deemed necessary.
 - f. Plans for criterion validity testing have been developed, and planning for this activity is now under way. See p. 49 for details.
- 12) The initial working version of the ONQI algorithm was finalized in July 2007.

This has been an excerpt from the ONQI Reference Manual. For the complete public version of the manual, please send an email to info@nuval.com, and it will be emailed to you.

Please note that due to the proprietary nature of the ONQI algorithm, some excerpts and details are omitted from the manual.

Thanks for your interest in ONQI and the NuVal Nutritional Scoring System.

BIBLIOGRAPHY

1. Department of Health and Human Services, Food and Drug Administration. Public Hearing on Use of Symbols to Communicate Nutrition Information, Consideration of Consumer Studies and Nutritional Criteria; September 10-11, 2007; Adelphi, MD. <http://www.cfsan.fda.gov/~comm/registre6.html>. Accessed October 15, 2007.
2. Kumpulainen JT. Chromium content of foods and diets. *Biol Trace Elem Res.* 1992;32:9-18.
3. US Department of Agriculture, Agricultural Research Service. USDA National Nutrient Database for Standard Reference, Release 18. 2005. <http://www.ars.usda.gov/Main/docs.htm?docid=15869>. Accessed January 8, 2006.
4. Institute of Medicine. *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids.* Washington, DC: National Academy Press; 2000. http://www.nap.edu/catalog.php?record_id=10026#toc. Accessed January 8, 2006.
5. Institute of Medicine. *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc;* 2000. http://books.nap.edu/openbook.php?record_id=10026&page=82. Accessed January 8, 2006.
6. Institute of Medicine. *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline;* 1998. http://books.nap.edu/openbook.php?record_id=6015&page=150. Accessed January 8, 2006.
7. Institute of Medicine. *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline;* 1998. http://books.nap.edu/openbook.php?record_id=6015&page=306. Accessed January 8, 2006.
8. Institute of Medicine. *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline;* 1998. http://books.nap.edu/openbook.php?record_id=6015&page=196. Accessed January 8, 2006.
9. Scott JM. Reduced folate status is common and increases disease risk. It can be corrected by daily ingestion of supplements or fortification. *Novartis Found Symp.* 2007;282:105-222.
10. Pitkin RM. Folate and neural tube defects. *Am J Clin Nutr.* 2007;85(1):285S-288S.
11. De Wals P, Tairou F, Van Allen MI, Uh SH, Lowry RB, Sibbald B, Evans JA, Van den Hof MC, Zimmer P, Crowley M, Fernandez B, Lee NS, Niyonsenga T. Reduction in neural-tube defects after folic acid fortification in Canada. *N Engl J Med.* 2007;12;357(2):135-142.

12. Green NS. Folic acid supplementation and prevention of birth defects. *J Nutr.* 2002;132(8 Suppl):2356S-2360S.
13. McKinley MC. Nutritional aspects and possible pathological mechanisms of hyperhomocysteinaemia: an independent risk factor for vascular disease. *Proc Nutr Soc.* 2000;59(2):221-237.
14. Dhonukshe-Rutten RA, de Vries JH, de Bree A, van der Put N, van Staveren WA, de Groot LC. Dietary intake and status of folate and vitamin B12 and their association with homocysteine and cardiovascular disease in European populations. *Eur J Clin Nutr.* 2007;Sep 12; [Epub ahead of print].
15. Institute of Medicine. *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*; 2000. http://books.nap.edu/openbook.php?record_id=9810&page=95. Accessed January 8, 2006.
16. Sasazuki S, Sasaki S, Tsubono Y, Okubo S, Hayashi M, Tsugane S. Effect of vitamin C on common cold: randomized controlled trial. *Eur J Clin Nutr.* 2006;60(1):9-17.
17. Li Y, Schellhorn HE. New developments and novel therapeutic perspectives for vitamin C. *J Nutr.* 2007;137(10):2171-2184.
18. Jacob RA, Sotoudeh G. Vitamin C function and status in chronic disease. *Nutr Clin Care.* 2002;5(2):66-74.
19. Institute of Medicine. *Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride*; 1997. http://books.nap.edu/openbook.php?record_id=5776&page=250. Accessed January 8, 2006.
20. Bischoff-Ferrari HA, Willett WC, Wong JB, Giovannucci E, Dietrich T, Dawson-Hughes B. Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *JAMA.* 2005;293(18):2257-2264.
21. Vieth R. The role of vitamin D in the prevention of osteoporosis. *Ann Med.* 2005;37(4):278-285.
22. Papadimitropoulos E, Wells G, Shea B, Gillespie W, Weaver B, Zytaruk N, Cranney A, Adachi J, Tugwell P, Josse R, Greenwood C, Guyatt G. Meta-analyses of therapies for postmenopausal osteoporosis. VIII: Meta-analysis of the efficacy of vitamin D treatment in preventing osteoporosis in postmenopausal women. *Endocr Rev.* 2002;23(4):560-569.
23. Pittas AG, Dawson-Hughes B, Li T, Van Dam RM, Willett WC, Manson JE, Hu FB. Vitamin D and calcium intake in relation to type 2 diabetes in women. *Diabetes Care.* 2006 Mar;29(3):650-656.

24. Holick MF. The vitamin D epidemic and its health consequences. *J Nutr.* 2005;135(11):2739S-2748S.
25. Cantorna MT, Mahon BD. Mounting evidence for vitamin D as an environmental factor affecting autoimmune disease prevalence. *Exp Biol Med (Maywood)*. 2004;229(11):1136-1142.
26. Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr.* 2004;80(6 Suppl):1678S-1688S.
27. Garland CF, Garland FC, Gorham ED, Lipkin M, Newmark H, Mohr SB, Holick MF. The role of vitamin D in cancer prevention. *Am J Public Health.* 2006;96(2):252-261.
28. Giovannucci E, Liu Y, Rimm EB, Hollis BW, Fuchs CS, Stampfer MJ, Willett WC. Prospective study of predictors of vitamin D status and cancer incidence and mortality in men. *J Natl Cancer Inst.* 2006;5;98(7):451-459.
29. Grant WB. Epidemiology of disease risks in relation to vitamin D insufficiency. *Prog Biophys Mol Biol.* 2006;92(1):65-79.
30. Skinner HG, Michaud DS, Giovannucci E, Willett WC, Colditz GA, Fuchs CS. Vitamin D intake and the risk for pancreatic cancer in two cohort studies. *Cancer Epidemiol Biomarkers Prev.* 2006;15(9):1688-1695.
31. Giovannucci E. The epidemiology of vitamin D and cancer incidence and mortality: a review (United States). *Cancer Causes Control.* 2005;16(2):83-95.
32. Gorham ED, Garland CF, Garland FC, Grant WB, Mohr SB, Lipkin M, Newmark HL, Giovannucci E, Wei M, Holick MF. Vitamin D and prevention of colorectal cancer. *J Steroid Biochem Mol Biol.* 2005;97(1-2):179-194.
33. Grant WB, Garland CF. A critical review of studies on vitamin D in relation to colorectal cancer. *Nutr Cancer.* 2004;48(2):115-123.
34. Institute of Medicine. *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*; 2000. http://books.nap.edu/openbook.php?record_id=9810&page=186. Accessed January 8, 2006.
35. Kontush K, Schekatolina S. Vitamin E in neurodegenerative disorders: Alzheimer's disease. *Ann N Y Acad Sci.* 2004;1031:249-262.
36. Traber MG. Heart disease and single-vitamin supplementation. *Am J Clin Nutr.* 2007;85(1):293S-299S.
37. Jackson RD, LaCroix AZ, Gass M, Wallace RB, Robbins J, Lewis CE, Bassford T, Beresford SA, Black HR, Blanchette P, Bonds DE, Brunner RL, Brzyski RG, Caan B, Cauley JA, Chlebowski

- RT, Cummings SR, Granek I, Hays J, Heiss G, Hendrix SL, Howard BV, Hsia J, Hubbell FA, Johnson KC, Judd H, Kotchen JM, Kuller LH, Langer RD, Lasser NL, Limacher MC, Ludlam S, Manson JE, Margolis KL, McGowan J, Ockene JK, O'Sullivan MJ, Phillips L, Prentice RL, Sarto GE, Stefanick ML, Van Horn L, Wactawski-Wende J, Whitlock E, Anderson GL, Assaf AR, Barad D (Women's Health Initiative Investigators). Calcium plus vitamin D supplementation and the risk of fractures. *N Engl J Med*. 2006;354(7):669-683.
38. Heaney RP, Weaver CM. Newer perspectives on calcium nutrition and bone quality. *J Am Coll Nutr*. 2005;24(6 Suppl):574S-581S.
39. Porthouse J, Cockayne S, King C, Saxon L, Steele E, Aspray T, Baverstock M, Birks Y, Dumville J, Francis R, Iglesias C, Puffer S, Sutcliffe A, Watt I, Torgerson DJ. Randomised controlled trial of calcium and supplementation with cholecalciferol (vitamin D3) for prevention of fractures in primary care. *BMJ*. 2005;330(7498):1003.
40. Shea B, Wells G, Cranney A, Zytaruk N, Robinson V, Griffith L, Hamel C, Ortiz Z, Peterson J, Adachi J, Tugwell P, Guyatt G. Calcium supplementation on bone loss in postmenopausal women. *Cochrane Database Syst Rev*. 2004;(1):CD004526.
41. Institute of Medicine. *Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride*; 1997. http://books.nap.edu/openbook.php?record_id=5776&page=71. Accessed January 8, 2006.
42. Ruidavets JB, Bongard V, Simon C, Dallongeville J, Ducimetière P, Arveiler D, Amouyel P, Bingham A, Ferrières J. Independent contribution of dairy products and calcium intake to blood pressure variations at a population level. *J Hypertens*. 2006;24(4):671-681.
43. Larsson SC, Bergkvist L, Rutegard J, Giovannucci E, Wolk A. Calcium and dairy food intakes are inversely associated with colorectal cancer risk in the Cohort of Swedish Men. *Am J Clin Nutr*. 2006 Mar;83(3):667-73; quiz 728-9.
44. Lin J, Zhang SM, Cook NR, Manson JE, Lee IM, Buring JE. Intakes of calcium and vitamin D and risk of colorectal cancer in women. *Am J Epidemiol*. 2005;161(8):755-764.
45. McCullough ML, Rodriguez C, Diver WR, Feigelson HS, Stevens VL, Thun MJ, Calle EE. Dairy, calcium, and vitamin D intake and postmenopausal breast cancer risk in the Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiol Biomarkers Prev*. 2005;(12):2898-2904.
46. Cho E, Smith-Warner SA, Spiegelman D, Beeson WL, van den Brandt PA, Colditz GA, Folsom AR, Fraser GE, Freudenheim JL, Giovannucci E, Goldbohm RA, Graham S, Miller AB, Pietinen P, Potter JD, Rohan TE, Terry P, Toniolo P, Virtanen MJ, Willett WC, Wolk A, Wu K,

- Yaun SS, Zeleniuch-Jacquotte A, Hunter DJ. Dairy foods, calcium, and colorectal cancer: a pooled analysis of 10 cohort studies. *J Natl Cancer Inst.* 2004;96(13):1015-1022.
47. Pittas AG, Dawson-Hughes B, Li T, Van Dam RM, Willett WC, Manson JE, Hu FB. Vitamin D and calcium intake in relation to type 2 diabetes in women. *Diabetes Care.* 2006;29(3):650-656.
48. Teegarden D. Calcium intake and reduction in weight or fat mass. *J Nutr.* 2003;(1):249S-251S.
49. Institute of Medicine. *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*; 2000. http://books.nap.edu/openbook.php?record_id=10026&page=290. Accessed January 8, 2006.
50. Institute of Medicine. *Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride*; 1997. http://books.nap.edu/openbook.php?record_id=5776&page=190. Accessed January 8, 2006.
51. Barbagallo M, Dominguez LJ, Resnick LM. Magnesium metabolism in hypertension and type 2 diabetes mellitus. *Am J Ther.* 2007;14(4):375-385.
52. He K, Song Y, Belin RJ, Chen Y. Magnesium intake and the metabolic syndrome: epidemiologic evidence to date. *J Cardiometab Syndr.* 2006;1(5):351-355.
53. Larsson SC, Wolk A. Magnesium intake and risk of type 2 diabetes: a meta-analysis. *J Intern Med.* 2007;262(2):208-214.
54. Pham PC, Pham PM, Pham SV, Miller JM, Pham PT. Hypomagnesemia in patients with type 2 diabetes. *Clin J Am Soc Nephrol.* 2007;2(2):366-373.
55. Institute of Medicine. *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate*; 2004. http://books.nap.edu/openbook.php?record_id=10925&page=186. Accessed January 8, 2006.
56. Institute of Medicine. *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*; 2000. http://books.nap.edu/openbook.php?record_id=10026&page=442. Accessed January 8, 2006.
57. Calder PC. n-3 polyunsaturated fatty acids, inflammation, and inflammatory diseases. *Am J Clin Nutr.* 2006;83(6 Suppl):1505S-1519S.
58. Robinson JG, Stone NJ. Antiatherosclerotic and antithrombotic effects of omega-3 fatty acids. *Am J Cardiol.* 2006;98(4A):39i-49i.
59. Wijendran V, Hayes KC. Dietary n-6 and n-3 fatty acid balance and cardiovascular health. *Annu Rev Nutr.* 2004;24:597-615.

60. Gebauer SK, Psota TL, Harris WS, Kris-Etherton PM. n-3 fatty acid dietary recommendations and food sources to achieve essentiality and cardiovascular benefits. *Am J Clin Nutr.* 2006;83(6 Suppl):1526S-1535S.
61. Lichtenstein AH, Appel LJ, Brands M, Carnethon M, Daniels S, Franch HA, Franklin B, Kris-Etherton P, Harris WS, Howard B, Karanja N, Lefevre M, Rudel L, Sacks F, Van Horn L, Winston M, Wylie-Rosett J. Diet and lifestyle recommendations revision 2006: a scientific statement from the American Heart Association Nutrition Committee. *Circulation.* 2006;114(1):82-96. .
62. Mozaffarian D, Rimm EB. Fish intake, contaminants, and human health: evaluating the risks and the benefits. *JAMA.* 2006;296(15):1885-1889.
63. Psota TL, Gebauer SK, Kris-Etherton P. Dietary omega-3 fatty acid intake and cardiovascular risk. *Am J Cardiol.* 2006;98(4A):3i-18i.
64. Bouzan C, Cohen JT, Connor WE, Kris-Etherton PM, Gray GM, Konig A, Lawrence RS, Savitz DA, Teutsch SM. A quantitative analysis of fish consumption and stroke risk. *Am J Prev Med.* 2005;29(4):347-352.
65. Harper CR, Jacobson TA. Usefulness of omega-3 fatty acids and the prevention of coronary heart disease. *Am J Cardiol.* 2005;96(11):1521-1529.
66. Konig A, Bouzan C, Cohen JT, Connor WE, Kris-Etherton PM, Gray GM, Lawrence RS, Savitz DA, Teutsch SM. A quantitative analysis of fish consumption and coronary heart disease mortality. *Am J Prev Med.* 2005;29(4):335-346.
67. Mozaffarian D. Does alpha-linolenic acid intake reduce the risk of coronary heart disease? A review of the evidence. *Altern Ther Health Med.* 2005;11(3):24-30
68. Covington MB. Omega-3 fatty acids. *Am Fam Physician.* 2004;70(1):133-140.
69. Kris-Etherton PM, Harris WS, Appel LJ; American Heart Association. Nutrition Committee. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation.* 2002;106(21):2747-2757.
70. Keenan JM, Goulson M, Shamliyan T, Knutson N, Kolberg L, Curry L. The effects of concentrated barley beta-glucan on blood lipids in a population of hypercholesterolaemic men and women. *Br J Nutr.* 2007;97(6):1162-1168.
71. Queenan KM, Stewart ML, Smith KN, Thomas W, Fulcher RG, Slavin JL. Concentrated oat beta-glucan, a fermentable fiber, lowers serum cholesterol in hypercholesterolemic adults in a randomized controlled trial. *Nutr J.* 2007;26:6:6.

72. Maki KC, Shinnick F, Seeley MA, Veith PE, Quinn LC, Hallissey PJ, Temer A, Davidson MH. Food products containing free tall oil-based phytosterols and oat beta-glucan lower serum total and LDL cholesterol in hypercholesterolemic adults. *J Nutr.* 2003;133(3):808-813.
73. Rose DJ, DeMeo MT, Keshavarzian A, Hamaker BR. Influence of dietary fiber on inflammatory bowel disease and colon cancer: importance of fermentation pattern. *Nutr Rev.* 2007;65(2):51-62.
74. Lin J, Zhang SM, Cook NR, Rexrode KM, Liu S, Manson JE, Lee IM, Buring JE. Dietary intakes of fruit, vegetables, and fiber, and risk of colorectal cancer in a prospective cohort of women (United States). *Cancer Causes Control.* 2005;16(3):225-233.
75. Kaline K, Bornstein SR, Bergmann A, Hauner H, Schwarz PE. The importance and effect of dietary fiber in diabetes prevention with particular consideration of whole grain products. *Horm Metab Res.* 2007;39(9):687-693.
76. Flight I, Clifton P. Cereal grains and legumes in the prevention of coronary heart disease and stroke: a review of the literature. *Eur J Clin Nutr.* 2006 Oct;60(10):1145-59.
77. Weickert MO, Möhlig M, Schöfl C, Arafat AM, Otto B, Viehoff H, Koebnick C, Kohl A, Spranger J, Pfeiffer AF. Cereal fiber improves whole-body insulin sensitivity in overweight and obese women. *Diabetes Care.* 2006;(4):775-780.
78. Jenkins DJ, Kendall CW, Vuksan V, Vidgen E, Parker T, Faulkner D, Mehling CC, Garsetti M, Testolin G, Cunnane SC, Ryan MA, Corey PN. Soluble fiber intake at a dose approved by the US Food and Drug Administration for a claim of health benefits: serum lipid risk factors for cardiovascular disease assessed in a randomized controlled crossover trial. *Am J Clin Nutr.* 2002;75(5):834-839.
79. Marlett JA, McBurney MI, Slavin JL; American Dietetic Association. Position of the American Dietetic Association: health implications of dietary fiber. *J Am Diet Assoc.* 2002;102(7):993-1000.
80. Neuhouser ML. Dietary flavonoids and cancer risk: evidence from human population studies. *Nutr Cancer.* 2004;50(1):1-7.
81. Graf BA, Milbury PE, Blumberg JB. Flavonols, flavones, flavanones, and human health: epidemiological evidence. *J Med Food.* 2005;8(3):281-290.
82. Yao LH, Jiang YM, Shi J, Tomas-Barberan FA, Datta N, Singanusong R, Chen SS. Flavonoids in food and their health benefits. *Plant Foods Hum Nutr.* 2004;59(3):113-122.
83. Ren MQ, Kuhn G, Wegner J, Chen J. Isoflavones, substances with multi-biological and clinical properties. *Eur J Nutr.* 2001;40(4):135-146.

84. Hodgson JM. Effects of tea and tea flavonoids on endothelial function and blood pressure: a brief review. *Clin Exp Pharmacol Physiol*. 2006;33(9):838-841.
85. Tamimi RM, Hankinson SE, Campos H, Spiegelman D, Zhang S, Colditz GA, Willett WC, Hunter DJ. Plasma carotenoids, retinol, and tocopherols and risk of breast cancer. *Am J Epidemiol*. 2005;161(2):153-160.
86. Holick CN, Michaud DS, Stolzenberg-Solomon R, Mayne ST, Pietinen P, Taylor PR, Virtamo J, Albanes D. Dietary carotenoids, serum beta-carotene, and retinol and risk of lung cancer in the alpha-tocopherol, beta-carotene cohort study. *Am J Epidemiol*. 2002; 156(6):536-547.
87. Mannisto S, Smith-Warner SA, Spiegelman D, Albanes D, Anderson K, van den Brandt PA, Cerhan JR, Colditz G, Feskanich D, Freudenheim JL, Giovannucci E, Goldbohm RA, Graham S, Miller AB, Rohan TE, Virtamo J, Willett WC, Hunter DJ. Dietary carotenoids and risk of lung cancer in a pooled analysis of seven cohort studies. *Cancer Epidemiol Biomarkers Prev*. 2004;13(1):40-48.
88. Institute of Medicine. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients)*;2005.
http://books.nap.edu/openbook.php?record_id=10490&page=542. Accessed January 8, 2006.
89. Connor WE, Connor SL. Dietary cholesterol and coronary heart disease. *Current Atherosclerosis Reports*. 2002;(4):425-432.
90. Mensink RP, Zock PL, Kester AD, Katan MB. Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 controlled trials. *Am J Clin Nutr*. 2003;77(5):1146-1155.
91. Boyd NF, Stone J, Vogt KN, Connelly BS, Martin LJ, Minkin S. Dietary fat and breast cancer risk revisited: a meta-analysis of the published literature. *Br J Cancer*. 2003; 89(9):1672-1685.
92. Mozaffarian D, Katan MB, Ascherio A, Stampfer MJ, Willett WC. Trans fatty acids and cardiovascular disease. *N Engl J Med*. 2006;354(15):1601-1613.
93. Willett WC. Trans fatty acids and cardiovascular disease-epidemiological data. *Atheroscler Suppl*. 2006;7(2):5-8.
94. Lopez-Garcia E, Schulze MB, Meigs JB, Manson JE, Rifai N, Stampfer MJ, Willett WC, Hu FB. Consumption of trans fatty acids is related to plasma biomarkers of inflammation and endothelial dysfunction. *J Nutr*. 2005;135(3):562-566.
95. Clifton PM, Keogh JB, Noakes M. Trans fatty acids in adipose tissue and the food supply are associated with myocardial infarction. *J Nutr*. 2004;134(4):874-879.

96. Riserus U. Trans fatty acids and insulin resistance. *Atheroscler Suppl.* 2006;7(2):37-39.
97. Odegaard AO, Pereira MA. Trans fatty acids, insulin resistance, and type 2 diabetes. *Nutr Rev.* 2006;64(8):364-372.
98. Bragulat E, de la Sierra A. Salt intake, endothelial dysfunction, and salt-sensitive hypertension. *J Clin Hypertens (Greenwich).* 2002;4(1):41-46.
99. Cohen HW, Alderman MH. Low sodium diet after DASH: has the situation changed? Dietary Approaches to Stop Hypertension. *Curr Hypertens Rep.* 2002;4(4):329-332.
100. Geleijnse JM, Kok FJ, Grobbee DE. Blood pressure response to changes in sodium and potassium intake: a meta-regression analysis of randomised trials. *J Hum Hypertens.* 2003;17(7):471-480.
101. Karppanen H, Mervaala E. Sodium intake and hypertension. *Prog Cardiovasc Dis.* 2006;49(2):59-75.
102. Bray GA, Nielsen SJ, Popkin BM. Consumption of high-fructose corn syrup in beverages may play a role in the epidemic of obesity. *Am J Clin Nutr.* 2004;79(4):537-543.
103. Malik VS, Schulze MB, Hu FB. Intake of sugar-sweetened beverages and weight gain: a systematic review. *Am J Clin Nutr.* 2006;84(2):274-288.
104. Schulze MB, Manson JE, Ludwig DS, Colditz GA, Stampfer MJ, Willett WC, Hu FB. Sugar-sweetened beverages, weight gain, and incidence of type 2 diabetes in young and middle-aged women. *JAMA.* 2004;292(8):927-934.
105. Vermunt SH, Pasma WJ, Schaafsma G, Kardinaal AF. Effects of sugar intake on body weight: a review. *Obes Rev.* 2003;4(2):91-99.
106. Institute of Medicine. *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*; 2000. http://books.nap.edu/openbook.php?record_id=9810&page=284. Accessed January 8, 2006.
107. Papp LV, Lu J, Holmgren A, Khanna KK. From selenium to selenoproteins: synthesis, identity, and their role in human health. *Antioxid Redox Signal.* 2007;9(7):775-806.
108. Lachance PA. Nutrient addition to foods: The public health impact in countries with rapidly westernizing diets. In: Bendich A, Deckelbaum RJ, eds. *Preventive Nutrition: The Comprehensive Guide for Health Professionals*. Totowa, NJ: Humana Press;1997.
109. WCRF/AICR (World Cancer Research Fund/American Institute for Cancer Research). *Food, Nutrition and the Prevention of Cancer: A Global Perspective*. Menasha, WI: BANTA Book Group;1997

110. Appel LJ, Sacks FM, Carey VJ, Obarzanek E, Swain JF, Miller ER 3rd, Conlin PR, Erlinger TP, Rosner BA, Laranjo NM, Charleston J, McCarron P, Bishop LM. Effects of protein, monounsaturated fat, and carbohydrate intake on blood pressure and serum lipids: results of the OmniHeart randomized trial. *JAMA*. 2005;294(19):2455-2464.
111. Rasmussen BM, Vessby B, Uusitupa M, Berglund L, Pedersen E, Riccardi G, Rivellese AA, Tapsell L, Hermansen K. Effects of dietary saturated, monounsaturated, and n-3 fatty acids on blood pressure in healthy subjects. *Am J Clin Nutr*. 2006;83(2):221-226.
112. Kris-Etherton PM. AHA science advisory: monounsaturated fatty acids and risk of cardiovascular disease. *J Nutr*. 1999;129(12):2280-2284.
113. Lada AT, Rudel LL. Dietary monounsaturated versus polyunsaturated fatty acids: which is really better for protection from coronary heart disease? *Curr Opin Lipidol*. 2003;14(1):41-46.
114. Kris-Etherton PM, Hecker KD, Binkoski AE. Polyunsaturated fatty acids and cardiovascular health. *Nutr Rev*. 2004;62(11):414-426.
115. Institute of Medicine. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients)*; 2005.
http://books.nap.edu/openbook.php?record_id=10490&page=589. Accessed January 8, 2006.
116. Ledikwe JH, Blanck HM, Khan LK, Serdula MK, Seymour JD, Tohill BC, Rolls BJ. Low-energy-density diets are associated with high diet quality in adults in the United States. *J Am Diet Assoc*. 2006;106(8):1172-1180.
117. Kral TV, Roe LS, Rolls BJ. Combined effects of energy density and portion size on energy intake in women. *Am J Clin Nutr*. 2004;79(6):962-968.
118. Drewnowski A. The role of energy density. *Lipids*. 2003;38(2):109-115.
119. Newby PK. Examining energy density: comments on diet quality, dietary advice, and the cost of healthful eating. *J Am Diet Assoc*. 2006;106(8):1166-1169.
120. Ledikwe JH, Blanck HM, Kettel Khan L, Serdula MK, Seymour JD, Tohill BC, Rolls BJ. Dietary energy density is associated with energy intake and weight status in US adults. *Am J Clin Nutr*. 2006;83(6):1362-1368.
121. Rolls BJ, Roe LS, Meengs JS. Reductions in portion size and energy density of foods are additive and lead to sustained decreases in energy intake. *Am J Clin Nutr*; 2006;83(1):11-17.
122. Rolls BJ, Drewnowski A, Ledikwe JH. Changing the energy density of the diet as a strategy for weight management. *J Am Diet Assoc*. 2005;105(5 Suppl 1):S98-103.

123. Ello-Martin JA, Ledikwe JH, Rolls BJ. The influence of food portion size and energy density on energy intake: implications for weight management. *Am J Clin Nutr.* 2005;82(1 Suppl):236S-241S.
124. Kendall CW, Augustin LS, Emam A, Josse AR, Saxena N, Jenkins DJ. The glycemic index: methodology and use. *Nestle Nutr Workshop Ser Clin Perform Programme.* 2006;11:43-56.
125. Dickinson S, Brand-Miller J. Glycemic index, postprandial glycemia and cardiovascular disease. *Curr Opin Lipidol.* 2005;16(1):69-75.
126. Schulze MB, Liu S, Rimm EB, Manson JE, Willett WC, Hu FB. Glycemic index, glycemic load, and dietary fiber intake and incidence of type 2 diabetes in younger and middle-aged women. *Am J Clin Nutr.* 2004;80(2):348-356.
127. Brand-Miller JC. Glycemic load and chronic disease. *Nutr Rev.* 2003;61(5 Pt 2):S49-55.
128. Liu S, Willett WC, Stampfer MJ, Hu FB, Franz M, Sampson L, Hennekens CH, Manson JE. A prospective study of dietary glycemic load, carbohydrate intake, and risk of coronary heart disease in US women. *Am J Clin Nutr.* 2000;71(6):1455-1461.
129. Augustin LS, Franceschi S, Jenkins DJ, Kendall CW, La Vecchia C. Glycemic index in chronic disease: a review. *Eur J Clin Nutr.* 2002;(11):1049-1071.
130. Willett W, Manson J, Liu S. Glycemic index, glycemic load, and risk of type 2 diabetes. *Am J Clin Nutr.* 2002;76(1):274S-280S.
131. Michaud DS, Liu S, Giovannucci E, Willett WC, Colditz GA, Fuchs CS. Dietary sugar, glycemic load, and pancreatic cancer risk in a prospective study. *J Natl Cancer Inst.* 2002;94(17):1293-1300.
132. Michaud DS, Fuchs CS, Liu S, Willett WC, Colditz GA, Giovannucci E. Dietary glycemic load, carbohydrate, sugar, and colorectal cancer risk in men and women. *Cancer Epidemiol Biomarkers Prev.* 2005;14(1):138-147.
133. Westrich BJ, Buzzard IM, Gatewood LC, McGovern PG. Accuracy and Efficiency of Estimating Nutrient Values in Commercial Food Products Using Mathematical Optimization. *J Food Comp Anal.* 1994;7: 223-239.
134. US Department of Agriculture, Agricultural Research Service. USDA National Nutrient Database for Standard Reference, Release 18. 2005. <http://www.nal.usda.gov/fnic/foodcomp>. Accessed October 1, 2007.

135. US Department of Agriculture, Agricultural Research Service. USDA Database for the Flavonoid Content of Selected Foods, Release 2.1. 2007.
<http://www.ars.usda.gov/Services/docs.htm?docid=6231> Accessed October 1, 2007.
136. Drewnowski A. Concept of a nutritious food: toward a nutrient density score. *Am J Clin Nutr.* 2005;82(4):721-732.